

Nontopological Finite Temperature Induced Fermion Number

I. J. R. Aitchison⁽¹⁾ and G. V. Dunne^(1,2)

⁽¹⁾ *Department of Theoretical Physics, Oxford University, 1 Keble Rd., Oxford OX1 3NP, U.K.*

⁽²⁾ *Research Centre for the Subatomic Structure of Matter, University of Adelaide, S.A. 5005, Australia*

We show that while the zero temperature induced fermion number in a chiral sigma model background depends only on the asymptotic values of the chiral field, at finite temperature the induced fermion number depends also on the detailed shape of the chiral background. We resum the leading low temperature terms to all orders in the derivative expansion, producing a simple result that can be interpreted physically as the different effect of the chiral background on virtual pairs of the Dirac sea and on the real particles of the thermal plasma. By contrast, for a kink background, not of sigma model form, the finite T induced fermion number is temperature dependent but topological.

The phenomenon of induced fermion number due to the interaction of fermions with topological backgrounds (e.g., solitons, vortices, monopoles, skyrmions) has many applications ranging from polymer physics to particle physics [1, 2, 3, 4, 5, 6, 7, 8]. The original fractional fermion number result of Jackiw-Rebbi [1] has a deep connection with the existence of spinless charged excitations in polymers [2]. The adiabatic analysis of Goldstone-Wilczek [3] in systems without conjugation symmetry has important implications for bag models [5], monopoles, and sigma models, which provide effective field theory descriptions of systems ranging from condensed matter, to AMO, to particle and nuclear physics [9]. The induced fermion number is related to the spectral asymmetry of the relevant Dirac operator, and mathematical results concerning index theorems [8] relate the fermion number to asymptotic topological properties of the background. At finite temperature, the situation is less clear. In several examples [10, 11, 12, 13, 14], the fermion number is known to be temperature dependent, but is still topological in the sense that the only dependence on the background field is through its asymptotic properties. In this Letter, we present a simple physical case for which this is not true : in a 1 + 1 dim chiral sigma model, the finite temperature induced fermion number depends on the detailed structure of the background. This contradicts a previous analysis [15] and claim [16] that the finite T fermion number is in general a topological quantity. We give a simple physical explanation of the origin of the nontopological dependence. Our analysis has been motivated in part by the results of [17] concerning the T dependence of anomalous amplitudes in nuclear decays.

Consider an abelian model in 1 + 1 dimensions with fermions interacting via scalar and pseudoscalar couplings to two bosonic fields ϕ_1 and ϕ_2 . For the purposes of this paper ϕ_1 and ϕ_2 can be considered as classical external fields. The Lagrangian is

$$\mathcal{L} = i \bar{\psi} \not{\partial} \psi - \bar{\psi} (\phi_1 + i \gamma_5 \phi_2) \psi \quad (1)$$

There are two especially interesting physical cases:

(i) kink case [1] :

$$\phi_1 = m \quad \text{and} \quad \phi_2(\pm\infty) = \pm \hat{\phi}_2 \quad (2)$$

(ii) sigma model case [3] :

$$\phi_1^2 + \phi_2^2 = m^2 \quad (3)$$

In the sigma model case (3), the interaction term in the Lagrangian (1) can be expressed as

$$m \bar{\psi} e^{i \gamma_5 \theta} \psi = m \bar{\psi} (\cos \theta + i \gamma_5 \sin \theta) \psi \quad (4)$$

At $T = 0$, both these cases have an induced topological current $J^\mu \equiv \langle \bar{\psi} \gamma^\mu \psi \rangle$ given by [3]

$$J^\mu = \frac{1}{2\pi} \epsilon^{\mu\nu} \partial_\nu \theta + \dots \quad (5)$$

where the angular field θ is defined by $\theta \equiv \arctan(\phi_2/\phi_1)$. The dots in (5) refer to higher derivative terms, which are all of the form of a total derivative of θ and its derivatives [4]. Thus, in particular, the induced fermion number, $N \equiv \int dx J^0$, is

$$N = \frac{1}{2\pi} \int_{-\infty}^{\infty} dx \theta' = \frac{1}{\pi} \hat{\theta} \quad (6)$$

where $\pm\hat{\theta}$ are the asymptotic values of $\theta(x)$ at $x = \pm\infty$. The fermion number N is topological as it depends only on $\hat{\theta}$, not on the detailed shape of $\theta(x)$. The conjugation symmetric case of Jackiw and Rebbi [1] is obtained by taking $m \rightarrow 0$ in the kink case (2), in which case $N \rightarrow \pm\frac{1}{2}$.

At nonzero temperature, the induced fermion number for a static background is [8, 14]

$$N = -\frac{1}{2} \int_{\mathcal{C}} \frac{dz}{2\pi i} \text{tr} \left(\frac{1}{H-z} \right) \tanh \left(\frac{\beta z}{2} \right) \quad (7)$$

where $\beta = 1/T$ is the inverse temperature, and $\text{tr}(\frac{1}{H-z})$ is the resolvent of the Dirac Hamiltonian H . The contour $\mathcal{C}$ is $(-\infty + i\epsilon, +\infty + i\epsilon)$ and $(+\infty - i\epsilon, -\infty - i\epsilon)$. By considering static backgrounds we avoid the well-known complications of finite temperature calculations in non-static backgrounds [18]. The technical part of the calculation of the induced fermion number (7) is the computation of the resolvent of H . Once this is done, the induced fermion number may be expressed as an integral representation, or as a sum by deforming the contour in (7) around the simple poles of the tanh function. For static backgrounds $\phi_1(x)$ and $\phi_2(x)$ in (1), the Dirac Hamiltonian is

$$H = -i\gamma^0\gamma^1\nabla + \gamma^0\phi_1(x) + i\gamma^0\gamma^5\phi_2(x) \quad (8)$$

where $\nabla \equiv \frac{d}{dx}$, and we will work with the Dirac matrices $\gamma^0 = \sigma_3$, $\gamma^1 = i\sigma_2$, and $\gamma^5 = -\sigma_1$. Also, note that only the *even* part (in terms of the argument z) of the resolvent $\text{tr}(\frac{1}{H-z})$ contributes to the induced fermion number N in (7). (This is most easily seen by deforming the contour around the poles of the tanh function.)

Consider first the kink case in (2). Then the even part of the resolvent can be computed exactly using a trace identity which is a special case of the Callias index theorem [8, 10, 19] (alternatively, it can be derived in a more elementary manner as an exact resummation of a SUSY derivative expansion [20]) :

$$\begin{aligned} \left[\text{tr} \left(\frac{1}{H-z} \right) \right]_{\text{even}} &= \text{tr} \left(\frac{m}{-(\nabla + \phi_2)(\nabla - \phi_2) + m^2 - z^2} \right) - \text{tr} \left(\frac{m}{-(\nabla - \phi_2)(\nabla + \phi_2) + m^2 - z^2} \right) \\ &= \frac{-m\hat{\phi}_2}{(m^2 - z^2) \sqrt{m^2 + \hat{\phi}_2^2 - z^2}} \end{aligned} \quad (9)$$

Then the induced fermion number (7) for the kink case (2) is

$$N = \frac{2}{\pi} \left(\frac{m\beta}{\pi} \right)^2 \sin\hat{\theta} \sum_{n=0}^{\infty} \frac{1}{((2n+1)^2 + (\frac{m\beta}{\pi})^2) \sqrt{(2n+1)^2 \cos^2\hat{\theta} + (\frac{m\beta}{\pi})^2}} \quad (10)$$




FIG. 1: Plots of πN , where N is the finite temperature fermion number (10) for the kink case (2), as a function of $\hat{\theta}$. These plots are for $m\beta/\pi$ taking values 0.5, 1, and 10, as labelled. As $T \rightarrow 0$, note that $\pi N \rightarrow \hat{\theta}$, as in (6).

where $\hat{\theta} \equiv \arctan(\hat{\phi}_2/m)$. This result is consistent with previous analyses [10], although these were much less explicit. The induced fermion number (10) is plotted in Fig. 1 as a function of $\hat{\theta}$ for various values of temperature. As $T \rightarrow 0$, this result reduces smoothly to the zero temperature result (6). Despite its complicated form, the nonzero temperature result (10) is still topological as it only refers to the background through $\hat{\theta}$.

In the sigma model case (3), the trace identity formulae (9) do not apply. Another approach is needed to evaluate the resolvent. One such approach is the derivative expansion [21], in which we assume that the spatial derivatives of the background fields are small compared to the fermion mass scale m . In other words, the backgrounds $\phi_1(x)$ and $\phi_2(x)$ are assumed to be slowly varying on the scale of the fermion Compton wavelength. Returning to the general Hamiltonian (8), the derivative expansion can be obtained by separating H^2 as

$$H^2 = \begin{pmatrix} -\nabla^2 + \phi_1^2 + \phi_2^2 & 0 \\ 0 & -\nabla^2 + \phi_1^2 + \phi_2^2 \end{pmatrix} + \begin{pmatrix} \phi_2' & i\phi_1' \\ -i\phi_1' & -\phi_2' \end{pmatrix} \quad (11)$$

and then expanding $\text{tr}\left(\frac{1}{H-z}\right) = \text{tr}\left((H+z)\frac{1}{H^2-z^2}\right)$ in powers of derivatives. A simple calculation to first order yields:

$$\left[\text{tr}\left(\frac{1}{H-z}\right)\right]_{\text{even}} = -\frac{1}{2} \int_{-\infty}^{\infty} dx \frac{(\phi_1 \phi_2' - \phi_2 \phi_1')}{(\phi_1^2 + \phi_2^2 - z^2)^{3/2}} + \dots \quad (12)$$

where the dots refer to terms with three or more derivatives.

In the kink case (2), where $\phi_1 = m$ is constant, this first order calculation actually reproduces the *exact* trace identity result (9). But in the sigma model case (3), where $\phi_1^2 + \phi_2^2 = m^2$ is a constant, the first order derivative expansion result (12) implies that:

$$\left[\text{tr}\left(\frac{1}{H-z}\right)\right]_{\text{even}} = -\frac{m^2}{2(m^2 - z^2)^{3/2}} \int_{-\infty}^{\infty} dx \theta' + \dots \quad (13)$$

So, to first order in the derivative expansion, the induced fermion number for the sigma model case is

$$N^{(1)} = \frac{1}{\pi} \left(\frac{m\beta}{\pi}\right)^2 \left(\sum_{n=0}^{\infty} \frac{1}{[(2n+1)^2 + (\frac{m\beta}{\pi})^2]^{3/2}}\right) \int_{-\infty}^{\infty} dx \theta' \quad (14)$$

which is simply the zero temperature answer (6) multiplied by a smooth function of T . As $T \rightarrow 0$, this prefactor reduces to $\frac{1}{2\pi}$, so the *full* zero temperature result (6) is regained. But at finite temperature, the first order (in the derivative expansion) formula (14) for the sigma model case differs from the kink case formula (10), even though each of (14) and (10) reduces to (6) at $T = 0$.

This raises the question of the higher order corrections to the derivative expansion (12). In the kink case (2), there are *no* higher order corrections to the even part of the resolvent in (11). This is due to the special form of the Hamiltonian in the kink background, which leads to the first order formula (12) agreeing with the exact trace identity result (9). There can, of course, be higher order corrections to the induced fermion number *density*, but these are all total (spatial) derivatives, and do not contribute to the integrated induced fermion number, even at nonzero temperature.

But in the sigma model case (3), where the trace identity does not apply, the situation is very different. Going to the next order in the derivative expansion, we find

$$\begin{aligned} \left[\text{tr}\left(\frac{1}{H-z}\right)\right]_{\text{even}} &= -\frac{m^2}{2(m^2 - z^2)^{3/2}} \int dx \theta' - \frac{m^2}{8(m^2 - z^2)^{5/2}} \int dx \theta''' \\ &\quad - \frac{m^2(4z^2 + m^2)}{16(m^2 - z^2)^{7/2}} \int dx (\theta')^3 + \dots \end{aligned} \quad (15)$$

where the dots refer to terms involving five or more derivatives. For a chiral background with $\theta(x)$ approaching its asymptotic values exponentially fast, the term $\int dx \theta'''$ vanishes. But $\int dx (\theta')^3$ does not vanish. Thus, the first order induced fermion number (14) acquires a third order correction:

$$N^{(3)} = \frac{\beta^2}{8\pi^3} \left(\frac{m\beta}{\pi}\right)^2 \left(\sum_{n=0}^{\infty} \frac{[-4(2n+1)^2 + (\frac{m\beta}{\pi})^2]}{[(2n+1)^2 + (\frac{m\beta}{\pi})^2]^{7/2}}\right) \int dx (\theta')^3 \quad (16)$$

This is **not** just a function of the asymptotic value $\hat{\theta}$ of the chiral field $\theta(x)$; it also depends on the actual *shape* of $\theta(x)$. Thus, the induced fermion number is no longer topological. This contradicts [15], where it is stated that the first order derivative expansion contribution (14) is the full answer. However, the energy trace prefactor in (16) vanishes at $T = 0$, so the nontopological third order contribution (16) vanishes at $T = 0$. Thus, the nontopological nature of the finite temperature induced fermion number is still consistent (at this order) with the topological nature of the zero temperature induced fermion number (6).

We now turn to a physical explanation of why, in the sigma model case, the finite temperature induced charge is more sensitive to the background field than at zero temperature. Note first of all that the chiral background acts like a static but spatially inhomogeneous electric field, as can be seen by making a local chiral rotation [4]: $\psi \rightarrow \tilde{\psi} = e^{i\theta\gamma_5/2}\psi$. In terms of these chirally rotated fields the Lagrangian (1), with interaction (4), becomes

$$\mathcal{L} = i\tilde{\psi}\not{\partial}\tilde{\psi} - m\tilde{\psi}\tilde{\psi} - \tilde{\psi}\gamma^0\frac{\theta'}{2}\tilde{\psi} \quad (17)$$

(The chiral rotation leads to an anomalous Jacobian in the path integral, but this does not affect the induced fermion number.) Thus, the chiral field acts as an inhomogeneous $A_0(x) = \frac{1}{2}\theta'(x)$, leading to an electric field

$$E(x) = \frac{1}{2}\theta''(x) \quad (18)$$

Given that $\theta(x)$ itself has a kink-like spatial profile, the electric field is such that it changes sign as a function of x , as shown in Fig. 2 (we choose $\theta' > 0$). This electric field acts on the Dirac sea to polarize the vacuum by aligning the virtual vacuum dipoles of the Dirac sea, producing a localized build-up of charge near the kink center. But at nonzero temperature, the electric field also has an effect on the thermal plasma, as we show below.

First, consider the full derivative expansion (12) of the even part of the resolvent, at low but nonzero temperature. At fifth order, there are three independent terms involving θ'''' , $\theta'''(\theta')^2$, and $(\theta')^5$. The θ'''' term vanishes when integrated over x , but the other two terms are generally nonzero. However, as $T \rightarrow 0$ the $(\theta')^5$ term dominates the $\theta'''(\theta')^2$ term. Indeed, for low temperature, the dominant term with $(2l-1)$ derivatives in the derivative expansion (12) involves $(\theta')^{2l-1}$. Then, using the chirally rotated form (17) of the Lagrangian, the dominant term at $(2l-1)^{\text{th}}$ order is simply:

$$N_{\text{dom}}^{(2l-1)} = \left(T \sum_{n=-\infty}^{\infty} \int \frac{dk}{2\pi} \frac{\text{tr}([\gamma^0(\not{p} + m)]^{2l})}{(p^2 + m^2)^{2l}} \right) \int dx \left(\frac{\theta'}{2} \right)^{2l-1} \quad (19)$$

with Euclidean $p = (\omega_n, k)$ and $\omega_n = (2n+1)\pi T$ the Matsubara modes.

At zero temperature, all these terms $N^{(2l-1)}$ vanish, except for $l = 1$. This fact is not obvious; it involves highly nontrivial cancellations between terms in the expansion of the trace. But at nonzero temperature, all the terms in




FIG. 2: For a kink-like chiral field $\theta(x)$ (dashed line), the corresponding electric field in (18) has the form shown in the solid line, producing a vacuum polarization charge distribution localized near the kink center, roughly following the dotted line θ' .

(19) are non-vanishing. Moreover, they have a remarkably simple low temperature ($T \ll m$) limit:

$$N^{(2l-1)} = \delta_{l,1} \int dx \frac{\theta'}{2\pi} - \sqrt{\frac{2mT}{\pi}} e^{-m/T} \frac{1}{(2l-1)!} \int dx \left(\frac{\theta'}{2T} \right)^{2l-1} + \dots \quad (20)$$

Thus, in the low temperature limit, we can resum the *entire* derivative expansion, to obtain the induced fermion number in the sigma model case (3,4) :

$$N = \int dx \frac{\theta'}{2\pi} - \sqrt{\frac{2mT}{\pi}} \int dx e^{-m/T} \sinh \left(\frac{\theta'}{2T} \right) + \dots \quad (21)$$

where the dots refer to subleading terms for $T \ll m$.

Several features of this result (21) deserve comment. First, note that at zero temperature, only the first term survives, producing the familiar result (6) that the induced fermion number depends on the chiral field $\theta(x)$ only through its asymptotic value $\hat{\theta} \equiv \theta(\infty)$. At zero temperature, one can invoke Lorentz invariance to constrain the form of higher order corrections to (5) to be total derivatives [4], but these arguments do not apply at finite temperature. We see this in (21): the temperature dependent corrections are not total derivatives of terms made from θ and its derivatives. At nonzero temperature this shows clearly that the induced fermion number is nontopological - it depends also on the detailed shape of $\theta(x)$. Second, the resummed exponential factors $e^{-(m \mp \theta'/2)/T}$ in (21) are consistent with the derivative expansion assumption that $\theta' \ll m$. Finally, the form of these exponential factors suggests an interpretation of the derivative expansion as an adiabatic change of the local Fermi level with a local chemical potential $\mu = -\theta'/2$, which once again is only sensible in the derivative expansion regime where $\theta' \ll m$.

To make this physical picture more precise, we can interpret the result (21) as follows. The first, topological, term refers to the induced charge coming from the polarization of the Dirac sea. This is temperature independent as the short-lived virtual “electron-positron dipoles” of the Dirac sea do not come to thermal equilibrium. The next term in (21) corresponds to the induced charge arising from the response of the real charges in the thermal plasma to the spatially inhomogeneous electric field (18). Indeed, the linear response [22] of the plasma at low temperature to such an electric field yields an induced fermion number density

$$\rho(x) = \int \frac{dk}{2\pi} f(x, k) \quad (22)$$

where the static distribution function $f(x, k)$ satisfies the Boltzmann equation

$$v \frac{\partial}{\partial x} f(x, k) = -E(x) \frac{\partial}{\partial k} f(x, k) \quad (23)$$

where $v = k/\sqrt{k^2 + m^2}$. Regarding $\mu = -\frac{1}{2}\theta'(x)$ as a local chemical potential, (23) is satisfied by local Fermi particle and antiparticle distribution functions

$$f_{\pm}(x, k) = \frac{1}{e^{\beta(\sqrt{k^2 + m^2} \mp \mu)} + 1} \quad (24)$$

Inserting $f = f_+ - f_-$ into (22), we obtain precisely the second, nontopological, term in (21) in the low temperature limit.

At $T=0$, the fermion number may be defined as a sharp observable [23]; but at $T > 0$, thermal fluctuations introduce an rms deviation. Thus, the finite T fermion number in (7,21) is a thermal expectation value $\langle N \rangle$, as in the monopole cases [12, 13, 14]. We have estimated $\langle N^2 \rangle - \langle N \rangle^2$, in the derivative expansion regime, in an analogous manner to the computation presented here for $\langle N \rangle$. We find that the rms deviation vanishes at $T=0$, but at nonzero T can be significant compared to the thermal shift in (21). Details of this will be reported elsewhere.

To conclude, we comment briefly on possible implications of these results for models in other dimensions for which there is an induced fermion number due to some nontrivial background. In $2+1$ dimensions, fermions in a static magnetic background acquire an induced charge that is topological, expressed in terms of the net magnetic flux of the background. At finite temperature, the induced charge remains topological, but is multiplied by a smooth function of the temperature [11]. In $3+1$ dimensions, fermions in a static Dirac monopole background acquire an induced charge that is temperature dependent at finite T , but still only depends on the background through the total magnetic charge and the self-adjoint extension parameter [12, 14]. A more interesting case is a static 't Hooft-Polyakov monopole background, which has a characteristic size scale. Consider, for example, the coupling

$$\mathcal{L}_{\text{int}} = \bar{\psi} (\not{A} + \phi + i\gamma_5 m) \psi \quad (25)$$

where ψ is an isodoublet fermion, A_μ is a static $SU(2)$ 't Hooft-Polyakov monopole, and ϕ is the corresponding static Higgs field. We have computed the finite temperature induced fermion number, using the $3+1$ trace identity used in the zero temperature case [24], and we find *precisely* the same expression (10) as in the $1+1$ kink case, with the identification $\hat{\theta} = \arctan(\hat{\phi}/m)$, where $\hat{\phi}$ is the asymptotic value of the magnitude $|\phi| = \sqrt{\phi^a \phi^a}$ of the Higgs field. Given that (10) reduces to (6) at $T = 0$, this monopole result is consistent with the familiar zero temperature result [3, 24] that the induced fermion number is proportional to $\hat{\theta}$ [25]. The $3+1$ dimensional analogue of the $1+1$ sigma model case (3,4) is the sigma model with coupling

$$\begin{aligned}\mathcal{L}_{\text{int}} &= m\bar{\psi}(\pi_0 + i\gamma_5 \vec{\pi} \cdot \vec{\tau})\psi \\ &= m\bar{\psi}\left(\frac{1}{2}(g + g^\dagger) + \frac{1}{2}(g - g^\dagger)\gamma_5\right)\psi\end{aligned}\quad (26)$$

where $\vec{\tau}$ are $su(2)$ generators, the fields π_0 and $\vec{\pi}$ are constrained by $\pi_0^2 + \vec{\pi}^2 = 1$, and the fields g in the second line are defined by $g = \pi_0 + i\vec{\pi} \cdot \vec{\tau}$. At zero temperature, there is an induced topological charge density

$$J^0 = \frac{1}{24\pi^2} \epsilon^{ijk} \text{tr} (g^{-1} \partial_i g g^{-1} \partial_j g g^{-1} \partial_k g) \quad (27)$$

The corresponding zero temperature integrated charge is given by the winding number of the background field g at zero temperature. We conjecture that at finite temperature this induced charge will acquire additional nontopological contributions similar to those found here for the $1+1$ sigma model case.

Acknowledgements:

This work has been supported in part (GD) by the U.S. Department of Energy grant DE-FG02-92ER40716.00, and by PPARC. GD thanks Balliol College, Oxford, for a Visiting Fellowship, and the Theoretical Physics Department at Oxford, and the CSSM at Adelaide for their hospitality and support.

-
- [1] R. Jackiw and C. Rebbi, “Solitons with Fermion Number 1/2”, Phys. Rev. D **13** (1976) 3398.
 - [2] W. Su, J. R. Schrieffer, A. Heeger, “Solitons in Polyacetylene”, Phys. Rev. Lett. **42** (1979) 1698, “Soliton Excitations in Polyacetylene”, Phys. Rev. B **22** (1980) 2099; M. Rice, “Charged II-Phase Kinks in Lightly Doped Polyacetylene”, Phys. Lett. A **71** (1979) 152.
 - [3] J. Goldstone and F. Wilczek, “Fractional Quantum Numbers on Solitons”, Phys. Rev. Lett. **47** (1981) 986.
 - [4] R. MacKenzie and F. Wilczek, “Illustrations of Vacuum Polarization by Solitons”, Phys. Rev. D **30** (1984) 2194, “Examples of Vacuum Polarization by Solitons”, Phys. Rev. D **30** (1984) 2260; F. Wilczek, “Adiabatic Methods in Field Theory”, in 1984 TASI Lectures in Elementary Particle Physics, Ann Arbor, D. Williams (Ed.).
 - [5] J. Goldstone, R. Jaffe, “Baryon Number in Chiral Bag Models”, Phys. Rev. Lett. **51** (1983) 1518.
 - [6] E. D’Hoker and E. Farhi, “Decoupling a Fermion whose Mass is generated by a Yukawa Coupling: The General Case”, Nucl. Phys. B **248** (1984) 59, “Decoupling a Fermion in the Standard Electro-Weak Theory”, Nucl. Phys. B **248** (1984) 77; E. D’Hoker and J. Goldstone, “The Derivative Expansion of the Fermion Number Current”, Phys. Lett. B **158** (1985) 429.
 - [7] D. Diakonov, V. Petrov, P. Pobylitsa, “A Chiral theory of Nucleons”, Nucl. Phys. B **306** (1988) 809; R. D. Ball, “Chiral Gauge Theory”, Phys. Rep. **182** (1989) 1.
 - [8] For a review, see: A. Niemi and G. Semenoff, “Fermion Number Fractionization in Quantum Field Theory”, Phys. Rep. **135** (1986) 99.
 - [9] S. Weinberg, *Quantum Theory of Fields II*, (Cambridge, 1996).
 - [10] A. Niemi and G. Semenoff, “Fractional Fermion Number at Finite Temperature”, Phys. Lett. B **135** (1984) 121; V. Soni and G. Baskaran, “Depletion of Fractional Fermion Number of a Soliton at Finite Chemical Potential and Temperature”, Phys. Rev. Lett. **53** (1984) 523; W. Keil, “Comment on Fractionally Charged Solitons at Finite Temperature and Chemical Potential”, Phys. Lett. A **106** (1984) 89;
 - [11] K. Babu, A. Das and P. Panigrahi, “Derivative Expansion and the Induced Chern-Simons Term at Finite Temperature in (2+1) Dimensions”, Phys. Rev. D **36** (1987) 3725; G. Dunne, “Aspects of Chern-Simons Theories”, in Les Houches 1998, *Topological Aspects of Low Dimensional Systems*, A. Comtet, F. David, Th. Jolicoeur and S. Ouvry (Eds), (Springer, 1999).
 - [12] C. Coriano and R. Parwani, “The Electric Charge of a Dirac Monopole at Nonzero Temperature”, Phys. Lett. B **363** (1995) 71.
 - [13] A. Goldhaber, R. Parwani and H. Singh, “On the Fractional Electric Charge of a Magnetic Monopole at Nonzero Temperature”, Phys. Lett. B **386** (1996) 207.

- [14] G. Dunne and J. Feinberg, “Finite Temperature Effective Action in Monopole Background”, Phys. Lett. B **477** (2000) 474.
- [15] S. Midorikawa, “Fractional Charge at Finite Temperature”, Prog. Theor. Phys. **69** (1983) 1831.
- [16] A. Niemi, “Topological Solitons in a Hot and Dense Fermi Gas”, Nucl. Phys. B **251** (1985) 155.
- [17] R. Pisarski, L. Trueman, M. Tytgat, “Anomalous Amplitudes in a Thermal Bath”, Prog. Theor. Phys. Suppl. **131** (1998) 427.
- [18] A. Das, *Finite Temperature Field Theory*, (World Scientific, Singapore, 1997).
- [19] C. Callias, “Index Theorems on Open Spaces”, Commun. Math. Phys. **62** (1978) 213.
- [20] G. Dunne, “Derivative Expansion and Soliton Masses”, Phys. Lett. B **467** (1999) 238.
- [21] I. J. R. Aitchison and C. Fraser, “Derivative Expansion of Fermion Determinants : Anomaly-induced Vertices, Goldstone-Wilczek Currents, and Skyrme Terms”, Phys. Rev. D **31** (1985) 2605.
- [22] M. Le Bellac, *Thermal Field Theory* (Cambridge, 1996), Section 6.4.
- [23] S. Kivelson and J. R. Schrieffer, “Fractional charge, a sharp quantum observable”, Phys. Rev. B **25** (1982) 6447; A. Goldhaber and S. Kivelson, “Local charge versus Aharonov-Bohm charge”, Phys. Lett. B **255** (1991) 445.
- [24] M. Paranjape and G. Semenoff, “Spectral Asymmetry, Trace Identities and the Fractional Fermion Number of Magnetic Monopoles”, Phys. Lett. B **132** (1983) 369.
- [25] This is similar to, but slightly different from, the point-monopole case considered in [13] where the theta parameter is related to the boundary conditions used in the point limit.